



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO9031



ADULT AMBULATORY INFUSION ORDER
Darbepoetin alfa (ARANESP)
Maintenance for Oncology Patients

Page 1 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

****This order set is for MAINTENANCE DOSING ONLY. Patients should have received first dose via the INITIATION order set with either chemotherapy-induced anemia or myelodysplastic syndrome selected as indications****

INDICATION: (Must check one)

- Chemotherapy-induced anemia
- Symptomatic anemia associated with myelodysplastic syndrome (MDS)

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. OHSU's formulary erythropoiesis stimulating agent (ESA) is darbepoetin alfa (ARANESP). All orders for epoetin alfa (PROCRIT) will be converted to darbepoetin alfa using equivalent therapeutic interchange dosing listed in the table below. Providers who prefer to use epoetin alfa must specify a reason for its use and utilize an alternate ordering form.
3. Patients receiving concurrent treatment with Iron Sucrose (VENOFER) and/or Vitamin B12 cannot receive ESA treatment on the same day. Patients may be on prophylactic oral iron supplementation concurrent with ESA treatment as long as supplementation for the prevention of iron deficiency is necessary due to ESA therapy alone.
4. Serum ferritin and transferrin saturation (TSAT) must be performed every month during initial (ESA) treatment and at least every 3 months during stable ESA treatment (serum ferritin >100 ng/mL, and TSAT >20%). Therapy with darbepoetin alfa may continue **only if hemoglobin DOES NOT exceed 11 g/dL.**
5. **For patients with chemotherapy-induced anemia:** The medical record must document the provider's rationale for determining the anemia is "chemotherapy-induced." Anemia must be secondary to myelosuppressive anticancer chemotherapy in solid tumors, multiple myeloma, lymphoma, or lymphocytic leukemia. Treatment should be limited to the 8 weeks following myelosuppressive chemotherapy.
6. **For patients with symptomatic anemia from MDS:** The patient must be symptomatic and his/her life expectancy must be >3 months. The medical record must display documentation that a bone marrow biopsy has been reviewed by a provider and is consistent with the diagnosis of MDS. The marrow blast count must be <5%



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Darbepoetin alfa (ARANESP)
Maintenance for Oncology Patients

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

LABS:

- Hemoglobin & Hematocrit, Routine, ONCE, **every visit** (every two or three weeks – prior to every dose)
- CMP, Routine, ONCE, every visit (every two or three weeks – prior to every dose)
- Ferritin (serum), Routine, ONCE, **every 12 weeks**
- Iron and TIBC (serum), Routine, ONCE, **every 12 weeks**
- Labs already drawn within _____ days – Labs scanned with orders

MEDICATIONS:

- **darbepoetin alfa (ARANESP), subcutaneous, ONCE**

Dose: (must check one)

- 150 mcg
- 200 mcg
- 300 mcg

Interval:

- Every (2 or 3) ____ weeks x ____ doses (4 or 6) for a total of 12 weeks of therapy

OTHER:

Conversion from epoetin alfa (PROCRIT) to darbepoetin alfa (ARANESP): Initial adult dosing

Epoetin alfa dose (units/week)	Darbepoetin alfa dose (mcg/week)
<1500	6.25
1500-2499	6.25
2500-4999	12.5
5000-10,999	25
11,000-17,999	40
18,000-33,999	60
34,000-89,999	100
≥90,000	200

In patients receiving epoetin alfa 2-3 times weekly, darbepoetin should be given once weekly. If epoetin is administered once weekly, darbepoetin should be given once every 2 weeks. Darbepoetin dosing every 2 weeks should be determined by adding the 2 weekly epoetin alfa doses, then convert to appropriate corresponding darbepoetin dose. Doses should be titrated to hemoglobin response thereafter

NURSING ORDERS:

1. Patients receiving concurrent treatment with Iron Sucrose (VENOFER) and/or Vitamin B12 cannot receive ESA treatment on the same day.
2. TREATMENT PARAMETERS – Hold treatment and call provider if hemoglobin is greater than 11, serum ferritin is less than or equal to 100 ng/mL, transferrin saturation is less than or equal to 20% or if blood pressure is greater than 180 systolic or 100 diastolic
3. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Darbepoetin alfa (ARANESP)
Maintenance for Oncology Patients

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Please check the appropriate box for the patient's preferred clinic location:



TUALITY HEALTHCARE
An OHSU Partner

Infusion Services
364 SE 8th Ave, Medical Plaza Suite 108B
Hillsboro, OR 97123
Phone number: (503) 681-4124
Fax number: (503) 681-4120



Celilo Cancer Center
1800 E 19th St
The Dalles, OR 97058
Phone number: (541) 296-7585
Fax number: (541) 296-7610